




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K	A2	(11) International Publication Number: WO 00/13648 (43) International Publication Date: 16 March 2000 (16.03.00)
(21) International Application Number: PCT/DK99/00640 (22) International Filing Date: 22 November 1999 (22.11.99) (30) Priority Data: PA 1999 00920 25 June 1999 (25.06.99) DK (71) Applicant (for all designated States except US): H. LUNDBECK A/S [DK/DK]; Ottiliavej 9, DK-2500 Valby-Copenhagen (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): PETERSEN, Hans [DK/DK]; Guldagervej 11, DK-2720 Vanløse (DK). ROCK, Michael, Harold [GB/DK]; Risbjergvej 28, DK-2650 Hvidovre (DK). SVANE, Henrik [DK/DK]; Skodsborgvej 12, DK-2830 Virum (DK). (74) Common Representative: H. LUNDBECK A/S; Att: John Meidahl Petersen, Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Upon the request of the applicant, before the expiration of the time limit referred to in Article 21(2)(a). Without international search report and to be republished upon receipt of that report. Without classification; title and abstract not checked by the International Searching Authority.</i>
(54) Title: METHOD FOR THE PREPARATION OF CITALOPRAM <div style="text-align: center;">  </div> <p style="text-align: right;">(IV)</p>		
(57) Abstract Method for the preparation of citalopram comprising reaction of a compound of Formula (IV) wherein R is halogen, or CF ₃ -(CF ₂) _n -SO ₂ -, n being 0 to 8, with a cyanide source in the presence of a palladium catalyst and a catalytic amount of Cu ⁺ or Zn ²⁺ , or with Zn(CN) ₂ in the presence of a palladium catalyst.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Method for the Preparation of Citalopram

The present invention relates to a method for the preparation of the well known antidepressant drug citalopram, 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofurancarbonitrile.

Background of the Invention.

Citalopram is a well known antidepressant drug that has now been on the market for some years and has the following structure:



Formula I


It is a selective, centrally acting serotonin (5-hydroxytryptamine; 5-HT) reuptake inhibitor, accordingly having antidepressant activities. The antidepressant activity of the compound has been reported in several publications, eg. J. Hyttel, *Prog. Neuro-Psychopharmacol. & Biol. Psychiat.*, **1982**, 6, 277-295 and A. Gravem, *Acta Psychiatr. Scand.*, **1987**, 75, 478-486. The compound has further been disclosed to show effects in the treatment of dementia and cerebrovascular disorders, EP-A 474580.

20

Citalopram was first disclosed in DE 2,657,271 corresponding to US 4,136,193. This patent publication describes the preparation of citalopram by one method and outlines a further method which may be used for preparing citalopram.

25 According to the process described, the corresponding 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofurancarbonitrile is reacted with 3-(N,N-dimethylamino)propyl-chloride in the presence of methylsulfinylmethide as condensing agent. The starting material was prepared from the corresponding 5-bromo derivative by reaction with cuprous cyanide.


30 According to the method, which is only outlined in general terms, citalopram may be obtained by ring closure of the compound:



Formula II

in the presence of a dehydrating agent and subsequent exchange of the 5-bromo group with cyano using cuprous cyanide. The starting material of Formula II is obtained from 5-bromophthalide by two successive Grignard reactions, i.e. with 4-fluorophenyl magnesium chloride and N,N-dimethylaminopropyl magnesium chloride, respectively.

A new and surprising method and an intermediate for the preparation of citalopram were described in US Patent No 4,650,884 according to which an intermediate of the formula



Formula III

is subjected to a ring closure reaction by dehydration with strong sulfuric acid in order to obtain citalopram. The intermediate of Formula III was prepared from 5-cyanophthalide by two successive Grignard reactions, i.e. with 4-fluorophenyl magnesium halogenide and N,N-dimethylaminopropyl magnesium halogenide, respectively.

15

Further processes are disclosed in International patent application Nos. WO 98019511, WO 98019512 and WO 98019513. WO 98019512 and WO 98019513 relate to methods wherein a 5-amino-, 5-carboxy- or 5-(sec. aminocarbonyl)phthalide is subjected to two successive Grignard reactions, ring closure and conversion of the resulting 1,3-dihydroisobenzofuran derivative to the corresponding 5-cyano compound, i.e. citalopram. International patent application No. WO 98019511 discloses a process for the manufacture of citalopram wherein a (4-substituted-2-hydroxymethylphenyl)-(4-fluorophenyl)methanol compound is subjected to ring closure and the resulting 5-substituted 1-(4-fluorophenyl)-1,3-dihydroisobenzofuran converted to the corresponding 5-cyano derivative which is alkylated with a (3-dimethylamino)propylhalogenide in order to obtain citalopram.


25

Finally, methods of preparing the individual enantiomers of citalopram are disclosed in US Patent No 4,943,590 from which it also appears that the ring closure of the intermediate of Formula III may be carried out via a labile ester with a base.

- 5 With respect to the above methods for the preparation of citalopram the process comprising exchange of the 5-bromo group with cyano proved not to be very convenient in commercial scale, since the yield was rather low, the product was impure and in particular that it was difficult to separate the resulting citalopram from the corresponding 5-bromo compound.
- 10 It has now been found that citalopram may be obtained in a high yield as a very pure product by a new catalytic process in which 5-cyano is exchanged for a 5-halogen or a 5-triflate group of 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuran thus avoiding the extensive work up of the old cyanide exchange process.

15 **Summary of the invention**


Accordingly, the present invention relates to a novel method for the preparation of citalopram comprising reaction of a compound of Formula IV



Formula IV

- 20 wherein R is iodo, bromo, chloro, or $\text{CF}_3\text{-(CF}_2\text{)}_n\text{-SO}_2\text{-}$ wherein n is an integer in the range 0-8, incl., with a cyanide source, for example KCN, NaCN or $(\text{R}'_4\text{N})\text{CN}$ where R'_4 indicates four groups which may be the same or different and are selected from hydrogen and straight chain or branched C_{1-6} alkyl, in the presence of a palladium catalyst and a catalytic amount of Cu^+
- 25 or Zn^{2+} , or with $\text{Zn}(\text{CN})_2$ in the presence a palladium catalyst, and isolation of the corresponding 5-cyano compound, i.e. citalopram

4



Formula I

as the base or a pharmaceutically acceptable salt thereof.

- 5 In another aspect, the present invention provides the novel intermediates of Formula IV wherein R is $\text{CF}_3\text{-(CF}_2\text{)}_n\text{-SO}_2\text{-}$ wherein n is an integer in the range 0-8 or R is iodo.

In a further aspect the invention relates to the above process in which the compound of Formula IV is the S-enantiomer.

10

In yet another aspect, the present invention relates to an antidepressant pharmaceutical composition comprising citalopram manufactured by the process of the invention.

- By the process of the invention citalopram is obtained as a pure product in high yield thus
 15 reducing costly purification processes. Furthermore, the reaction may be carried out in more convenient solvents, at a low temperature and at a low excess of CN^- compared to the known cyano exchange process. The process has environmental advantages in that it only uses small amounts of heavy. Finally, this process gives an improved crystalline product enabling easy conversion to desired salts. The intermediates of Formula IV wherein R is $\text{CF}_3\text{-(CF}_2\text{)}_n\text{-SO}_2\text{-}$
 20 wherein n is an integer in the range 0-8 or R is iodo have been found to show pharmacological activity, i.e. 5-HT reuptake inhibiting effects, and accordingly they are useful as antidepressants

- The cyanide source used may be any useful source. Preferred sources are KCN, NaCN or
 25 $(\text{R}'_4\text{N})\text{CN}$ where R'_4 is as defined above. The cyanide source is used in a stoichiometric amount or in excess, preferably 1-2 equivalents are used per equivalent starting material of Formula IV. $\text{R}'_4\text{N}^+$ may conveniently be $(\text{Bu})_4\text{N}^+$. The cyanide compound is preferably NaCN or KCN or $\text{Zn}(\text{CN})_2$.

- 30 The palladium catalyst may be any suitable Pd(0) or Pd(II) containing catalyst, such as $\text{Pd}(\text{PPh}_3)_4$, $\text{Pd}_2(\text{dba})_3$, $\text{Pd}(\text{PPh})_2\text{Cl}_2$, etc. The Pd catalyst is conveniently used in an amount of 1-10, preferably 2-6, most preferably about 4-5 mol%.

Catalytic amounts of Cu^+ and Zn^{2+} , respectively, means substoichiometric amounts such as 0.1 - 5, preferably 1 - 3 eq. %. Conveniently, about $\frac{1}{2}$ eq. is used per eq. Pd. Any convenient source of Cu^+ and Zn^{2+} may be used. Cu^+ is preferably used in the form of CuI and Zn^{2+} is conveniently used as the $\text{Zn}(\text{CN})_2$ salt.

5

In a preferred embodiment of the invention, R is $\text{CF}_3-(\text{CF}_2)_n-\text{SO}_2-$ wherein n is an integer from the range 0 to 8 or R is bromo or iodo, most preferably $\text{CF}_3-(\text{CF}_2)_8-\text{SO}_2-$, CF_3-SO_2- , bromo or iodo, in particular bromo.

- 10 In another particularly preferred embodiment the compound of Formula IV is reacted with ZnCl_2 in the presence of a Palladium catalyst, preferably $\text{Pd}(\text{PPh}_3)_4$ (tetrakis(triphenylphosphine)palladium).

The intermediate of Formula IV wherein R is bromo or chloro may be prepared from bromo- and chlorophthalide, respectively, as described in DE 2,657,271 and the corresponding US 4,136,193. The iodo may be prepared analogously from the corresponding phthalide derivatives and the compounds wherein R is $\text{CF}_3-(\text{CF}_2)_n-\text{SO}_2-$ may be prepared from the corresponding hydroxy compounds by a conventional triflation reaction.

- 20 The reaction may be performed in any convenient solvent, preferably acetonitril, propionitrile, THF and ethylacetate.

Other reaction conditions, solvents, etc. are conventional conditions for such reactions and may easily be determined by a person skilled in the art.

25

The compound of general Formula I may be used as the free base or as a pharmaceutically acceptable acid addition salt thereof. As acid addition salts, such salts formed with organic or inorganic acids may be used. Exemplary of such organic salts are those with maleic, fumaric, benzoic, ascorbic, succinic, oxalic, bismethylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzene sulfonic and theophylline acetic acids, as well as the 8-halothephyllines, for example 8-bromothephylline. Exemplary of such inorganic salts are those with hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric and nitric acids.

35

The acid addition salts of the compounds may be prepared by methods known in the art. The base is reacted with either the calculated amount of acid in a water miscible solvent, such as acetone or ethanol, with subsequent isolation of the salt by concentration and cooling, or with

an excess of the acid in a water immiscible solvent, such as ethylether, ethylacetate or dichloromethane, with the salt separating spontaneously.

The pharmaceutical compositions of the invention may be administered in any suitable way and in any suitable form, for example orally in the form of tablets, capsules, powders or syrups, or parenterally in the form of usual sterile solutions for injection.

The pharmaceutical formulations of the invention may be prepared by conventional methods in the art. For example, tablets may be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: Corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvant or additive colourings, aroma, preservatives etc. may be used provided that they are compatible with the active ingredients.

Solutions for injections may be prepared by solving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to the desired volume, sterilisation of the solution and filling in suitable ampoules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

Examples

The invention is further illustrated by the following examples.

Example 1

Citalopram oxalate

Method 1

A mixture of $\text{Zn}(\text{CN})_2$ (1.2g, 0.01mol) and 1-(4'-fluorophenyl)-1-(3-dimethylaminopropyl)-5-bromophthalane (6.0g, 0.016mol) in DMF (40mL) was stirred at room temperature under an atmosphere of argon for 30 minutes. Dissolved oxygen was removed by bubbling argon through the reaction mixture for 10 minutes and then tetrakis(triphenylphosphine)palladium (0) (0.8g, 0.0007mol, 4.3mol%) was added. Then the reaction mixture was heated at 75 °C for 3 hrs, poured into water (200mL) and extracted with diethyl ether (2 x 100mL), dried (MgSO_4), filtered and concentrated under reduced pressure. The residue was dissolved in acetone (10mL) and a solution of oxalic acid (0.145g, 0.016mol) in acetone (10 mL) was added with stirring. The citalopram oxalate was isolated by filtration, washed with cold diethyl ether and dried in vacuo to pure citalopram, oxalate (6.1 g, 92 %)

Method 2

A mixture of 1-(4'-fluorophenyl)-1-(3-dimethylaminopropyl)-5-bromophthalane (2.5g, 0.007mol), NaCN (0.68 g, 0.014mol), and Zn(CN)₂ (0.014g, 0.00012mol) in THF (40 mL) were stirred at room temperature under an atmosphere of argon for 30 minutes. Then dissolved oxygen was removed by bubbling argon through the reaction mixture before the addition of tetrakis(triphenylphosphine)palladium (0) (0.3 g, 0.0003 mol, 3.7 mol%). Then the reaction mixture was heated at reflux overnight, cooled, diluted with diethyl ether, and then filtered through celite. The filtrate was washed with brine, dried (MgSO₄) and concentrated under reduced pressure. The residue was dissolved in acetone (50mL) and a solution of oxalic acid (0.63g, 0.007mol) in acetone (10 mL) was added with stirring. The Citalopram oxalate was isolated by filtration, washed with cold diethyl ether and dried in vacuo to pure citalopram, oxalate (2.4g, 82%)

Example 2

1-(4'-fluorophenyl)-1-(3-dimethylaminopropyl)-5-iodophthalane, oxalate.

A solution of 4-fluorophenylmagnesium bromide, prepared from 4-fluorobromobenzene (19.3 g, 0.11 mole) and magnesium turnings (2.92 g, 0.12 mol) in dry THF (100 mL), is added dropwise to a suspension of 5-iodophthalide (26.0 g, 0.1 mole) in dry THF (100 mL). The temperature is kept below 0 °C. After the addition is complete, the reaction mixture is stirred for 3 hours at 0 °C.

A second Grignard solution prepared from 3-dimethylaminopropyl chloride (14.6 g, 0.12 mole) and magnesium turnings (3.2 g, 0.13 mole) in dry THF (100 mL) is added to the reaction mixture. The temperature is kept below 0 °C during the addition. After the addition is complete the cooling is removed and the reaction mixture is stirred for an additional 2 hours at ambient temperature.

The reaction mixture is then poured into a mixture of ice water (200 mL) and a saturated solution of NH₄Cl (100 mL). THF is evaporated in vacuo. Toluene (200 mL) is added and the organic phase is separated and extracted with 1 M HCl (1 x 100 mL). The pH of the water phase is then adjusted to 9 by addition of 25 % NH₄OH (15 mL) and toluene (100 mL) is added. The reaction is left overnight at room temperature.

The organic phase is separated and 70 % sulfuric acid (10 mL) is added at room temperature. The reaction mixture is stirred at room temperature for 2 hours to complete the ring closure. 25 % NH₄OH (20 mL) is added and the organic phase is separated, filtered and evaporated in vacuo to give the crude title compound as its free base.

A sample of the crude material (5.0 g, 11.3 mmol) is dissolved in ethyl acetate and filtered through silica. Eluent 1: Ethyl acetate which is discarded. Eluent 2: Ethyl acetate:Triethyl

amine, 95:5 which is collected and evaporated in vacuo to give the title compound (3.5 g, 8.2 mmol) as its free base.

The oxalate salt is precipitated from acetone.

DSC onset: 82 °C and 195 °C. ¹H NMR (DMSO d-6, 250 MHz): 1.3 - 1.65 (2H,m), 2.15
5 (2H,t, J=10 Hz), 2.63 (6H,s), 2.87 (2H,t, J=10 Hz), 5.0-5.2 (2H, 2d, J= 12.5 Hz), 6.5 -
7.05 (2H,s (broad)), 7.16 (2H,t, J=7.5 Hz), 7.35 (1H,d, J=8.5 Hz), 7.55 (2H,dt, J= 1.2
Hz, J=7.5 Hz), 7.64 (1H,d, J=8.5 Hz), 7.69 (1H,s).

Example 3

10 *1-(3-Dimethylamino-1-propyl)-1-(4-fluorophenyl)-5-hydroxy-1,3-dihydroisobenzofurane,*
oxalate

A solution of 4-fluorophenylmagnesium bromide, prepared from 4-fluorobromobenzene
(24,0 g, 0,14 mole) and magnesium turnings (4,38 g, 0,17 mole) in dry THF (80 mL), is
15 added dropwise to a suspension of 5-hydroxyphthalide (10,0 g, 0,07 mole) in dry THF (100
mL) at a temperature below 8°C. The reaction mixture is stirred at room temperature
overnight after the addition is finished.

A second Grignard solution prepared from 3-dimethylaminopropyl chloride (8,50 g, 0,07
mole) and magnesium turnings (1,93 g, 0,07 mole) in dry THF (40 mL) and added to the
20 reaction mixture while the temperature is kept below 10°C. The reaction is left stirred
overnight.

The reaction mixture is poured into ice water (200 mL) and pH is adjusted to 7 with
ammonium chloride water (300 mL) resulting in separation of two phases. The water phase is
extracted with ethylacetate (300 mL) and then made basic to pH 8 - 9 with 25%(w/v)
25 ammonium hydroxide. The water phase is extracted with toluene/ethylacetate (3:2, 3x100
mL). The toluene extract is dried over anhydrous sodium sulphate and stirred with charcoal.
After filtration the solvent is evaporated in vacuo and the title compound is obtained as a oil
(10,2 g, 48%).

5,1 grams (16 mmol) of the obtained oil is dissolved in acetone (25 mL) and treated with
30 anhydrous oxalic acid (1,46 g, 0,016 mole). The mixture is left in the freezer overnight and
the precipitated oxalate is filtered off. Yield: 4,77 g

DSC onset 168°C. ¹H NMR (DMSO-d₆, 500 MHz): 1,36 - 1,58 (2H, m), 2,05 - 2,18 (2H, m),
2,63 (6H, s), 2,96 (2H, t, J = 6,5 Hz), 4,95 (1H, d, J = 12,5 Hz), 5,08 (1H, d, J = 12,5 Hz),
6,65 (1H, s), 6,70 (1H, d, J = 8,5 Hz), 7,14 (2H, t, J = 7,5 Hz), 7,24 (1H, d, J = 8,5 Hz) 7,52
35 (2H, dt, J = 7,5 J = 1,2 Hz), 9 - 10 (2H, broad s).

Anal. calc. for C₂₁H₂₄N₁F₁O₆: C, 62,20; H, 5,98; N, 3,46.

Found: C, 62,02; H, 5,97; N, 3,42.

Example 4

1-(3-Dimethylamino-1-propyl)-1-(4-fluorophenyl)-5-[(trifluoromethyl)sulfonyl-oxy]-1,3-dihydroisobenzofurane, oxalate.

- 5 1-(3-Dimethylamino-1-propyl)-1-(4-fluorophenyl)-5-hydroxy-1,3-dihydroisobenzofurane (1,79 g, 5,7 mmol) is dissolved in dichloromethane (35 ml) and cooled in ice/water bath. Under nitrogen trifluoromethane sulfonic acid chloride (0,73 ml, 6,8 mmol) is added dropwise keeping the temperature below 5°C. The reaction mixture is allowed to warm to room temperature overnight. Water (40 mL) and triethylamine (1 mL) are added and the
- 10 phases are separated. The water phase is extracted with dichloromethane (25 mL). The combined organic phases are dried over magnesium sulphate and the solvent evaporated in vacuo. The residue (2,09 grams of the title compound as its free base) is dissolved in acetone (10 mL) and treated with anhydrous oxalic acid (0,51 g, 5,7 mmol). After stirring at room temperature overnight the precipitate is filtered off. Yield: 0,84 g, 33%.
- 15 DSC onset 144°C. ¹H NMR (DMSO-d₆, 500 MHz): 1,37 - 1,57 (2H, m), 2,15 - 2,25 (2H, m), 2,61 (6H, s), 2,95 (2H, t, J = 9,4 Hz), 5,12 (1H, d, J = 12,5 Hz), 5,22 (1H, d, J = 12,5 Hz), 7,17 (2H, t, J = 6,3 Hz), 7,42 (1H, d, J = 7,8 Hz), 7,48 (1H, s), 7,59 (2H, dt, J = 6,3 Hz J = 1,2 Hz), 7,70 (1H, d, J = 7,8 Hz).
- Anal. calc. for C₂₂H₂₃N₁F₄O₈S₁: C, 49,16; H, 4,32; N, 2,61.
- 20 Found: C, 49,43; H, 4,36; N, 2,57.

Example 5


Citalopram, oxalate, Method 3

- 25 1-(3-Dimethylamino-1-propyl)-1-(4-fluorophenyl)-5-[(trifluoromethyl)sulfonyl-oxy]-1,3-dihydroisobenzofurane (1,02 g, 2,3 mmol), sodium cyanide (0,22 g, 4,6 mmol), copper iodide (0,05 g, 0,3 mmol) and tetrakis(triphenylphosphine)palladium(0) (0,125 g, 0,1 mmol) are suspended in acetonitrile (10 mL). The suspension is heated at reflux for 5 hours and then
- 30 allowed to cool to room temperature overnight with intensive stirring. Ethylacetate (30 mL) is added and the mixture is filtrated on celite. The filtrate is washed with brine (60 mL) and dried over magnesium sulphate before the solvent is removed in vacuo. The crude product is eluted on silica (eluent: ethylacetate, ethanol, triethylamine 75:25:4). Yield: 0,22 g, 30%.
- The oxalate salt is precipitated from acetone.

35


CLAIMS

1. A method for the preparation of citalopram comprising reaction of a compound of Formula IV



Formula IV

wherein R is halogen, or $\text{CF}_3\text{-(CF}_2\text{)}_n\text{-SO}_2\text{-}$ wherein n is an integer in the range 0-8, incl., with a cyanide source in the presence of a palladium catalyst and a catalytic amount of Cu^+ or Zn^{2+} , or with Zn(CN)_2 in the presence a palladium catalyst, and isolation of the corresponding 5-cyano compound, i.e. citalopram



Formula I

as the base or a pharmaceutically acceptable salt thereof.

2. The method of Claim 1, wherein the cyanide source is KCN, NaCN or $(\text{R}'_4\text{N})\text{CN}$ where R'_4 indicates four groups which may be the same or different and are selected from hydrogen and straight chain or branched C_{1-6} alkyl,

3. The method of Claim 1 or 2 wherein R is $\text{CF}_3\text{-(CF}_2\text{)}_n\text{-SO}_2\text{-}$ wherein n is an integer from the range 0 to 8, preferably $\text{CF}_3\text{-SO}_2\text{-}$.

4. The method of Claim 1, 2 or 3 wherein R is bromo or iodo.

5. The method of any of Claims 1 - 4 wherein the compound of Formula IV is reacted with ZnCl_2 in the presence of a Palladium catalyst, preferably $\text{Pd(PPh}_3)_4$.

5 6. The method of any of Claims 1 - 5 wherein the cyanide compound used is NaCN, KCN or Zn(CN)_2 .

7. The method of any of Claims 1 - 4 and 6 wherein the palladium catalyst is $\text{Pd(PPh}_3)_4$, $\text{Pd}_2(\text{dba})_3$ or $\text{Pd(PPh)}_2\text{Cl}_2$.

10

8. The method of Claim 7 wherein the palladium catalyst is $\text{Pd(PPh}_3)_4$.


9. The method of any of Claims 1 - 8 wherein the reaction is carried out in the presence of a catalytic amount of Cu^+ , preferably in the form of CuI.

15

10. The method of any of Claims 1 - 8 wherein the reaction is carried out in the presence of a catalytic amount of Zn^{2+} , preferably as Zn(CN)_2 .

11. A compound of Formula IV

20



Formula IV


wherein R is $\text{CF}_3\text{-(CF}_2)_n\text{-SO}_2\text{-}$ wherein n is an integer in the range 0-8 or R is iodo.

25 12. The method of any of Claims 1 - 10 wherein the compound of Formula IV is the S-enantiomer.

13. An antidepressant pharmaceutical composition comprising citalopram manufactured by the process of any of Claims 1 - 10.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 307/87	A3	(11) International Publication Number: WO 00/13648 (43) International Publication Date: 16 March 2000 (16.03.00)
(21) International Application Number: PCT/DK99/00640 (22) International Filing Date: 22 November 1999 (22.11.99) (30) Priority Data: PA 1999 00920 25 June 1999 (25.06.99) DK (71) Applicant (for all designated States except US): H. LUNDBECK A/S [DK/DK]; Ottiliavej 9, DK-2500 Valby-Copenhagen (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): PETERSEN, Hans [DK/DK]; Guldagervej 11, DK-2720 Vanløse (DK). ROCK, Michael, Harold [GB/DK]; Risbjergvej 28, DK-2650 Hvidovre (DK). SVANE, Henrik [DK/DK]; Skodsborgvej 12, DK-2830 Virum (DK). (74) Common Representative: H. LUNDBECK A/S; Att: John Meidahl Petersen, Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Upon the request of the application, before the expiration of the time limit referred to in Article 21(2)(a).</i> (88) Date of publication of the international search report: 13 July 2000 (13.07.00)
(54) Title: METHOD FOR THE PREPARATION OF CITALOPRAM <div style="text-align: center;">  <p>(IV)</p> </div> (57) Abstract <p>Method for the preparation of citalopram comprising reaction of a compound of Formula (IV) wherein R is halogen, or CF₃-(CF₂)_n-SO₂-, n being 0 to 8, with a cyanide source in the presence of a palladium catalyst and a catalytic amount of Cu⁺ or Zn²⁺, or with Zn(CN)₂ in the presence of a palladium catalyst.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	IS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00640

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 307/87

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4136193 A (KLAUS P. BØGESØ ET AL), 23 January 1979 (23.01.79)	1-12
X	-- -----	13

☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 April 2000

Date of mailing of the international search report

03 -05- 2000

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson/ELY

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/DK 99/00640

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4136193 A	23/01/79	AT 359488 B	10/11/80
		AT 360001 B	10/12/80
		AT 360002 B	10/12/80
		AT 571979 A	15/05/80
		AT 572079 A	15/05/80
		AT 947276 A	15/04/80
		AU 509445 B	15/05/80
		AU 2107377 A	13/07/78
		BE 850401 A	14/07/77
		CA 1094087 A	20/01/81
		CH 626886 A	15/12/81
		CH 632258 A	30/09/82
		CH 632259 A	30/09/82
		DE 2657013 A,C	28/07/77
		DK 13177 A	15/07/77
		ES 454980 A	01/04/78
		FI 63754 B,C	29/04/83
		FI 770073 A	15/07/77
		FR 2338271 A,B	12/08/77
		GB 1526331 A	27/09/78
		IE 44055 B	29/07/81
		JP 1368581 C	11/03/87
		JP 52105162 A	03/09/77
		JP 61035986 B	15/08/86
		NL 192451 B,C	01/04/97
		NL 7700244 A	18/07/77
		NO 147243 B	22/11/82
		NO 770109 A	15/07/77
		NZ 183001 A	02/06/78
		SE 429551 B,C	12/09/83
		SE 7614201 A	15/07/77
		ZA 7700057 A	30/11/77



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K	A2	(11) International Publication Number: WO 00/13648
		(43) International Publication Date: 16 March 2000 (16.03.00)

(21) International Application Number: PCT/DK99/00640

(22) International Filing Date: 22 November 1999 (22.11.99)

(30) Priority Data:

PA 1999 00920 25 June 1999 (25.06.99) DK

(71) Applicant (for all designated States except US): H. LUNDBECK A/S [DK/DK]; Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PETERSEN, Hans [DK/DK]; Guldagervej 11, DK-2720 Vanløse (DK). ROCK, Michael, Harold [GB/DK]; Risbjergvej 28, DK-2650 Hvidovre (DK). SVANE, Henrik [DK/DK]; Skodsborgvej 12, DK-2830 Virum (DK).


(74) Common Representative: H. LUNDBECK A/S; Att: John Meidahl Petersen, Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

*Upon the request of the applicant, before the expiration of the time limit referred to in Article 21(2)(a).**Without international search report and to be republished upon receipt of that report.**Without classification; title and abstract not checked by the International Searching Authority.*

(54) Title: METHOD FOR THE PREPARATION OF CITALOPRAM



(IV)

(57) Abstract

Method for the preparation of citalopram comprising reaction of a compound of Formula (IV) wherein R is halogen, or CF_3 -(CF_2)_n- SO_2 -, n being 0 to 8, with a cyanide source in the presence of a palladium catalyst and a catalytic amount of Cu^+ or Zn^{2+} , or with $\text{Zn}(\text{CN})_2$ in the presence of a palladium catalyst.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 March 2000 (16.03.2000)

PCT

(10) International Publication Number
WO 00/13648 A3

(51) International Patent Classification⁷: C07D 307/87

(21) International Application Number: PCT/DK99/00640

(22) International Filing Date:
22 November 1999 (22.11.1999)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 1999 00920 25 June 1999 (25.06.1999) DK

(71) Applicant (for all designated States except US): H. LUNDBECK A/S [DK/DK]; Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PETERSEN, Hans [DK/DK]; Guldagervej 11, DK-2720 Vanløse (DK). ROCK, Michael, Harold [GB/DK]; Risbjergvej 28, DK-2650 Hvidovre (DK). SVANE, Henrik [DK/DK]; Skodsborgvej 12, DK-2830 Virum (DK).

(74) Common Representative: H. LUNDBECK A/S; Att: John Meidahl Petersen, Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,

ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- With amended claims.
- Upon request of the applicant, before the expiration of the time limit referred to in Article 21(2)(a).

(88) Date of publication of the international search report:
13 July 2000

Date of publication of the amended claims: 15 March 2001


(15) Information about Correction:

Previous Correction:

see PCT Gazette No. 19/2000 of 11 May 2000, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR THE PREPARATION OF CITALOPRAM




(57) Abstract: Method for the preparation of citalopram comprising reaction of a compound of Formula (IV) wherein R is halogen, or CF₃-(CF₂)_n-SO₂-, n being 0 to 8, with a cyanide source in the presence of a palladium catalyst and a catalytic amount of Cu⁺ or Zn²⁺, or with Zn(CN)₂ in the presence of a palladium catalyst.

WO 00/13648 A3

AMENDED CLAIMS


[received by the International Bureau on 25 October 2000 (25.10.00);
original claims 1, 3, 11 and 12 amended; remaining claims unchanged (2 pages)]

1. A method for the preparation of citalopram comprising reaction of a compound of
Formula IV



Formula IV

- wherein R is halogen, or $\text{CF}_3\text{-(CF}_2\text{)}_n\text{-SO}_2\text{-O-}$ wherein n is an integer in the range 0-8, incl.,
with a cyanide source in the presence of a palladium catalyst and a catalytic amount of Cu^+ or
 Zn^{2+} , or with Zn(CN)_2 in the presence a palladium catalyst, and isolation of the corresponding
5-cyano compound, i.e. citalopram



Formula I

as the base or a pharmaceutically acceptable salt thereof.

2. The method of Claim 1, wherein the cyanide source is KCN, NaCN or $(\text{R}'_4\text{N})\text{CN}$
where R'_4 indicates four groups which may be the same or different and are selected from
hydrogen and straight chain or branched C_{1-6} alkyl,

3. The method of Claim 1 or 2 wherein R is $\text{CF}_3\text{-(CF}_2\text{)}_n\text{-SO}_2\text{-O-}$ wherein n is an integer
from the range 0 to 8, preferably $\text{CF}_3\text{-SO}_2\text{-O-}$.

4. The method of Claim 1, 2 or 3 wherein R is bromo or iodo.

5. The method of any of Claims 1 - 4 wherein the compound of Formula IV is reacted with ZnCl_2 in the presence of a palladium catalyst, preferably $\text{Pd(PPh}_3)_4$.

5 6. The method of any of Claims 1 - 5 wherein the cyanide compound used is NaCN, KCN or Zn(CN)_2 .

7. The method of any of Claims 1 - 4 and 6 wherein the palladium catalyst is $\text{Pd(PPh}_3)_4$, $\text{Pd}_2(\text{dba})_3$ or $\text{Pd(PPh)}_2\text{Cl}_2$.

10

8. The method of Claim 7 wherein the palladium catalyst is $\text{Pd(PPh}_3)_4$.


9. The method of any of Claims 1 - 8 wherein the reaction is carried out in the presence of a catalytic amount of Cu^+ , preferably in the form of CuI.

15

10. The method of any of Claims 1 - 8 wherein the reaction is carried out in the presence of a catalytic amount of Zn^{2+} , preferably as Zn(CN)_2 .

11. A compound of Formula IV

20



Formula IV

wherein R is $\text{CF}_3-(\text{CF}_2)_n-\text{SO}_2-\text{O}-$ wherein n is an integer in the range 0-8 or R is iodo.

25 12. The method of any of Claims 1 - 10 wherein the compound of Formula IV is the S-enantiomer.

13. An antidepressant pharmaceutical composition comprising citalopram manufactured by the process of any of Claims 1 - 10.

30